

Competition of Branches

M. B. Hastings

CNLS, MS B258, Los Alamos National Laboratory, Los Alamos, NM 87545, hastings@cnls.lanl.gov
(May 16, 2001)

We consider a general model of branch competition that automatically leads to a critical branching configuration. This model is inspired by the $4 - \eta$ expansion of the dielectric breakdown model (DBM), but the mechanism of arriving at the critical point may be of relevance to other branching systems as well, such as fractures. The exact solution of this model clarifies the direct renormalization procedure used for the DBM, and demonstrates nonperturbatively the existence of additional irrelevant operators with complex scaling dimensions leading to discrete scale invariance. The anomalous exponents are shown to depend upon the details of branch interaction; we contrast with the branched growth model (BGM) in which these exponents are universal to lowest order in $1 - \nu$, and show that the BGM includes an inherent branch interaction different from that found in the DBM. We consider stationary and non-stationary regimes, corresponding to different growth geometries in the dielectric-breakdown model.

Diffusion-limited aggregation [1] produces complicated fractal structures by branch competition. Recently, by considering the dielectric breakdown model [2] (DBM), a controlled renormalization group was developed within a $4 - \eta$ expansion for this class of models [3]. This expansion is based on considering an aggregate as a collection of strictly one-dimensional branches; at $\eta = 4$, the aggregate consists of a single branch. Given that the probability of a single branch pair surviving for time t is proportional to $1/t^\nu$, to use the terminology of the branched growth model [4], a direct renormalization group was developed in an expansion in $1 - \nu = (4 - \eta)/2$. It was found that, without fine tuning, the system arrives at a critical point characterized by a scale-invariant tip-splitting rate so that the renormalized probability of a branch surviving for a time t is proportional to $1/t$ and the rate of branch production and branch death balance at all scales. However, branching structures are common in other physical systems, and a similar $1/t$ kernel has been found in fracture systems [5].

Thus, we will consider a more general model of branch competition, inspired by the $4 - \eta$ renormalization group. This model will be exactly solvable, clarifying the direct renormalization procedure employed for the DBM. We will find that the exponents depend on the bare kernel and on the details of branch interaction, becoming trivial as the $\nu \rightarrow 1$. Recently [6], it was shown that within the branched growth model (BGM) the fractal dimension depends only on ν for small ν ; here, this is shown not to be true for general models of branch interaction, though the fact that the numerical values of the exponents within the $4 - \eta$ [3] and branched growth [6] expansions are similar is an indication that the BGM is a useful approximation. However, we show that the physics of the BGM involves certain assumptions which do not hold for the DBM, so that the BGM result is not exact for the DBM. The exact solution of the general model will also reveal the presence of additional, irrelevant operators with complex scaling dimension, indicating the presence of discrete scale invariance, which has been argued to exist in DLA [7].

Finally, we will consider this model within two separate regimes. One regime will be analogous to the steady state regime in the cylindrical geometry in the DBM, while the other will be a non-stationary state, which corresponds to an initial condition of a single infinitely long vertical branch in cylindrical geometry, over time scales much shorter than the time required for the aggregate to reach the scale of the cylinder. We will find that the exponents characterizing local fractal dimension and mass-radius scaling are the same. This indicates that the difference between these dimensions observed in radial geometry [8], as well as the non-trivial affine exponents observed in the early stages of growth in cylindrical geometry [9] from initial conditions consisting of a horizontal line, are due not just to the non-stationary nature of the growth but also to the different geometry and initial conditions. Another motivation to consider different geometries is to compare the branched growth model computation, performed in a non-stationary regime, to the $4 - \eta$ RG computation, performed in a stationary regime.

A Model of Branch Competition — The dynamical state of the system at a given time t will be defined by a binary tree, with a set of times t_i , one for each branch point in the tree, defining the time at which that branch pair was produced. There are two dynamical processes, tip-splitting which leads to the production of additional branches and branch competition. The first process is accounted for by assigning a rate g at which each tip splits, changing the topology of the tree by adding an additional branch point at that tip with time $t_i = t$. See Fig. 1.

The second process is accounted for by assigning a probability of a branch point i being removed from the tree due to competition of branches. When the branch point is created, one of the two branches below the branch point is randomly designated as the weaker branch, and the other as the stronger. When the branch point is removed, the weaker branch is removed from the tree. See Fig. 2. We pick the probability of removing branch point i to be

$$\nu \theta(t - t_i - 1) \frac{1}{t - t_i}, \quad (1)$$

plus additional corrections due to branch interaction. To include the effects of branch interaction, we pick a phenomenological model for the interaction. We pick a scale factor $x < 1$, and declare that if on the weaker branch there exists another branch point with branching time exactly equal to $t - x(t - t_i)$ then the weaker branch will be removed. In this case, the lower branch point will have lived for a time x shorter than the branch point i .

This model has the essential features of the $4 - \eta$ renormalization group. One difference is that within that procedure the details of branch competition had to be determined numerically by integrating the trajectories of several competing branches, where here they may be determined phenomenologically by the scale factor x . Another difference has to do with the time scale. Within the DBM, each tip has a given growth measure and the tip-splitting rate and probability of removing branch points are proportional to the growth measure at the tip normalized by the total growth measure for the cluster. Here, we instead take all tips to have the same growth measure (so that all tips have the same probability of splitting) and we do not normalize the tip-splitting rate. The lack of normalization simply changes the overall time scale; the choice of the same growth measure for all tips is simply taken to make the model more tractable analytically and does not alter the essential physics.

In addition, in the $4 - \eta$ renormalization group, branch points may also be removed as a result of interactions with parent branches of similar scale, rather than just as a result of daughter branches. We will consider this possibility later.

Survival Probability — Let a branch pair be created at time $t = 0$. Define $s(t)$ to be the probability that the branch pair survives until time t , assuming that the branch pair is not destroyed by removal of one of its parents. We find

$$\partial_t s(t) = -\frac{\nu \theta(t-1)s(t)}{t} - g(1-x)s(t)s(xt), \quad (2)$$

as $g(1-x)s(xt)dt$ is the probability of creating a branch point at time $(1-x)t$ which survives until time t . The growth rules have been chosen so that Eq. (2) is exact; the process of creating and then removing a branch point does not effect the distribution of branches on the remaining stronger branch below that branch point.

Searching for a scaling solution, suppose $gs(t) = A/t$. We find $1 - \nu = (1-x)A/x$. The constant A sets the probability, in time t , of producing a branch which remains in the tree for time t ; this is the dimensionless (scale invariant) tip-splitting rate [3]. As $\nu \rightarrow 1$, $A \rightarrow 0$.

This result can also be obtained by a direct renormalization procedure [3] in which one expands the survival probability in powers of $1 - \nu$ and g : to zeroth order in g , $gs(t) = A/t + A(1-\nu) \log t/t + \mathcal{O}(1-\nu)^2$. Then, to first order in g we find $gs(t) = A/t + A(1-\nu) \log t/t - A^2(1-x) \log t/(xt)$, so that a fixed point of ts is reached only for the given value of A .

To investigate the approach to the scaling solution, suppose instead $gs(t) = \frac{A}{t}(1 + f(t))$. Linearizing Eq. (2) about $f = 0$, we find

$$\frac{\partial f(t)}{\partial \log t} = -(1-\nu)f(xt). \quad (3)$$

Eq. (3) is translationally invariant in $\tau = \log t$, and has solutions $f = e^{k\tau}$, where the eigenvalue k is the scaling dimension. We find

$$k = -(1-\nu)x^k. \quad (4)$$

When $\nu \approx 1$, the eigenvalue with largest real part is $k = -(1-\nu) + \mathcal{O}(1-\nu)^2$. Then, for $\nu \approx 1$, where this eigenvalue is small we can make an approximation that $f(xt) = f(t)$ and approximate the nonlinear problem by $\partial_t s(t) = -\nu s(t)/t - g(1-x)s^2(t)/x$, which can be solved exactly for $s(t)$, $t > 1$ as $s(t) = (1-\nu)/(g(1-x)t/x + ct^\nu)$, where c is an arbitrary constant.

For $1 - \nu$ small, Eq. (4) has two solutions for real k , as well as an infinity of solutions with complex k . All these k have negative real part and describe irrelevant perturbations. For $x^{1-\nu} = e^{-1/e}$ the two real solutions merge, and for $x^{1-\nu} < e^{-1/e}$ all solutions have complex k . The presence of complex eigenvalues indicates that $s(t)$ has an oscillatory behavior and that there is a discrete scale invariance in the corrections to scaling. The scale of this discrete scale invariance is in general *not* equal to x , so that it is not simply an artifact of the particularly simple form of branch interaction chosen, but rather a result of the fact that branches separated by a finite range of scales interact. As $x^{1-\nu}$ is decreased, one finds a k with vanishing real part when $k = i\pi/(2 \log x)$, so $x^{1-\nu} = e^{-\pi/2}$. For $x^{1-\nu} < e^{-\pi/2}$, there are complex eigenvalues with positive real part which describe *relevant* perturbations. In this case, we have found numerically that $f(t)$ is driven to a new fixed point with non-decaying log-periodic oscillations, with scale not in general equal to x . The model considered here is useful for analyzing these effects, which are beyond perturbation theory in $1 - \nu$.

Consider correlations: the probability, given that there is a branch point at time t_1 remaining in the tree until time t'_1 , that there is another branch point at time t_2 remaining in the tree until time t'_2 . To study these, generalize the survival probability to a function $s(t, t')$, the probability that a branch point created at time t survives until time $t + t'$. We obtain the equation

$$\partial_{t'} s(t, t') = -\frac{\nu \theta(t' - 1) s(t, t')}{t'} - g(1 - x) s(t, t') s((1 - x)t + xt', xt'). \quad (5)$$

Defining $gs(t, t') = \frac{A}{t}(1 + f(t, t'))$ and linearizing we find

$$\partial_{t'} f(t, t') = -\frac{(1 - \nu)}{t'} f(t + (1 - x)t', xt'). \quad (6)$$

This equation is translationally invariant in t and so we look for solutions $f(t, t') = e^{ilt} g(t')$. We find $\partial_{t'} g(t') = -\frac{(1 - \nu)}{t'} g(xt') e^{ilxt'}$. For $l = 0$, this is the same as Eq. (4). Generally, for $t' \ll l^{-1}$, we find the same discrete scale invariant solution as for Eq. (4) as above. For $t' \gg l^{-1}$, the perturbations $f(t, t')$ decay more rapidly due to the oscillations of the exponential. Thus, the two time scales t, t' are related, and we find discrete scale invariance in both.

Different Geometries— We now consider the local “fractal dimension” of the cluster. The length of a branch is defined to be $t - t_i$, where t_i is the time at which the pair to which that branch belongs was created. The largest such $t - t_i$ sets a length scale for the cluster. The mass of a branch is defined to be the length of that branch, plus the sum, over all branch points j which lie within that branch at time t , of $t - t_j$, plus the sum, over all branch points which were added to that branch and later removed, of the elapsed time between the addition and removal of the branch point.

Define $m(t)$ to be the average mass of a branch which is removed from the tree after a time t has elapsed since its creation (hence, this is a weaker branch). By averaging over the weights of sidebranches, we obtain the exact recursion relation

$$m(t) = t + g \int_0^t dt' \int_1^{xt'} dt'' m(t'') (-\partial_{t''} s(t'')). \quad (7)$$

Assuming a power law $m(t) = t^D$, one finds $\alpha(\alpha - 1) = Ax^{\alpha-1}$. For $\nu \approx 1$, we find

$$D = 1 + (1 - \nu) \frac{x}{1 - x} + \mathcal{O}(1 - \nu)^2. \quad (8)$$

Consider two different geometries. If we start the tree with a single branch and follow the dynamics above, this is analogous to starting the DBM with a single branch as a seed configuration and letting the cluster grow. In this case, Eq. (8) provides a scaling of the mass with the time. In another geometry, analogous to the cylindrical geometry, we modify the dynamics to always remove a weaker branch if $t - t_i > T$, for some T setting a scale. In this case, after an initial non-stationary regime lasting for a time of order T , the mass of the cluster increases linearly with t , with a rate proportional to T^{D-1} : the mass of the largest branches, times the probability of producing such a branch. Thus, within this model the mass-radius scaling and local fractal dimension are the same up to a trivial difference of unity. We can generalize the model by including a possibility of removing a branch due to the presence of a parent branch of comparable size. Then, the probability of the earliest branch point i surviving till time t will scale as $(t - t_i)^{-a} t_i^{-b}$ where $a + b = 1$. One will again find that the average time required to produce a branch point surviving for time t is of order t and the mass-radius scaling and local fractal dimension will again be equivalent.

Comparison to Branched Growth Model— It has been shown [6] that there exists a $1 - \nu$ expansion for the BGM similar to the $4 - \eta$ expansion for the DBM. One elegant feature of this expansion is that the lowest order fractal dimension is obtained without considering interaction of branches, but simply from the bare constant ν . This seems surprising, as we have found within the model above, and within the $4 - \eta$ expansion for the DBM, that unless we include branch interaction, the tip-splitting rate grows at large scales for $\nu < 1$, and a scale invariant fixed point is not reached.

The resolution of this is that the BGM is defined in a way which inherently includes effects of branch competition. In the BGM, a branch is assumed to have a probability $1/m^\nu$ of surviving until it reaches a mass m , while within the model above the probability is defined in terms of the probability to survive for a time t . Now, given that a scale invariant fixed point can only be reached if a branch has probability $1/t$ of surviving for time t , then we must have the relation that $1/m^\nu = 1/t$, so that $m = t^{1/\nu}$, and the fractal dimension is $1/\nu$.

More formally, $m(t) = t + At \log t$ to order $(1 - \nu)^0, A^1$. We do not have an exact Eq. (7) for the BGM, but this equation is still correct to lowest order. One may still define a survival probability $s(t)$ for a branch within the

BGM, and $\partial_t s(t) = -\nu \partial_t m(t) s(t) / m(t)$ so $s(t) = 1/t + (1 - \nu) \log t / t - A \log t / t$ to order $(1 - \nu)^1, A^1$, and we find $A = 1 - \nu = D - 1$.

However, this particular form of branch competition in the BGM is not that found in the DBM, as found by numerically following the evolution of three branches. In some circumstances production of a daughter branch can actually reduce the competition of the parent branch with its sister [3]; it is only when summing over all configurations that the increase in branch competition is obtained. Further, the competition of branches which is inherent in the BGM involves only an increased competition of a branch pair due to daughter branches. However, within the $4 - \eta$ expansion, it was necessary to consider the interaction of branches with daughter and parent branches (which is in fact the strongest interaction numerically) to obtain the correct result. Thus, both the present toy model and the BGM are approximations to the physics near $\eta = 4$, although the BGM dynamics serves also as a good approximation for $\eta = 1$ [4].

Conclusion— We have examined a simple, solvable model for branching, finding a fixed point with a scale invariant tip splitting rate. The direct RG for the model is exact at lowest order. The DBM has similar behavior and a similar perturbative RG. The exact solution enables us nonperturbatively to find additional irrelevant operators leading to a discrete scale invariance. We contrast the behavior of this model, and the DBM, with that of the BGM, for which a similar $1 - \nu$ expansion is available.

Acknowledgements— I thank T. C. Halsey for discussions. This work was supported by DOE grant W-7405-ENG-36.

- [1] T. A. Witten and L. M. Sander, Phys. Rev. Lett. **47**, 1400 (1981).
- [2] L. Niemeyer, L. Pietronero, and H. J. Wiesmann, Phys. Rev. Lett. **52**, 1033 (1984).
- [3] M. B. Hastings, preprint cond-mat/0104344.
- [4] T. C. Halsey and M. Leibig, Phys. Rev. A **46**, 7793 (1992).
- [5] J. P. Bouchaud et. al., Phys. Rev. Lett. **71**, 2240 (1993).
- [6] T. C. Halsey, preprint cond-mat/0105047.
- [7] D. Sornette et. al., Phys. Rev. Lett. **76**, 251 (1996).
- [8] B. B. Mandelbrot et. al., Europhys. Lett. **29**, 599 (1995).
- [9] C. Evertsz, Phys. Rev. A **42**, 1830 (1990).




FIG. 1. Creation of a new branch point.




FIG. 2. Removal of a branch below the circled branch point.